arXiv:1201.5733v2 [math.DS] 21 May 2012

ON THE SELF-SIMILARITY PROBLEM FOR GAUSSIAN-KRONECKER FLOWS

K. FRĄCZEK, J. KUŁAGA, AND M. LEMAŃCZYK

Abstract. It is shown that a countable symmetric multiplicative subgroup $G = -H \cup H$ with $H \subset \mathbb{R}_+^*$ is the group of self-similarities of a Gaussian-Kronecker flow if and only if H is additively $\mathbb{Q}$-independent. In particular, a real number $s \neq \pm 1$ is a scale of self-similarity of a Gaussian-Kronecker flow if and only if s is transcendental. We also show that each countable symmetric subgroup of $\mathbb{R}^*$ can be realized as the group of self-similarities of a simple spectrum Gaussian flow having the Foiaş-Stratila property.

Introduction

Assume that $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ is a (measurable) measure-preserving flow acting on a standard probability Borel space $(X, \mathcal{B}, \mu)$. Given $s \in \mathbb{R}^*$, one says that it is a *scale of self-similarity* of $\mathcal{T}$ if $\mathcal{T}$ is isomorphic to $\mathcal{T}_s := (T_{st})_{t\in\mathbb{R}}$. Denote by $I(\mathcal{T})$ the set of all scales of self-similarities of $\mathcal{T}$. Then $\mathcal{T}$ is called *self-similar* if $I(\mathcal{T}) \neq \{\pm 1\}$. Classical examples of self-similar flows are given by horocycle flows where $I(\mathcal{T})$ equals either $\mathbb{R}^*$ or $\mathbb{R}_+^*$ [19]. A systematic study of the problem of self-similarity has been done recently in [4] and [6]. In particular, $I(\mathcal{T})$ turns out to be a multiplicative subgroup of $\mathbb{R}^*$ ([6]) which is Borel ([4]), and one of the main problems in this domain is to classify all Borel subgroups of $\mathbb{R}^*$ that may appear as groups of self-similarities of ergodic flows; see also [3], [13], [24], [25] for a recent contribution to other aspects of the problem of self-similarity of ergodic flows. From this point of view the subclass of so called GAG flows [17][1] of the class of Gaussian flows is especially attractive since self-similarities appear there as natural invariants, see (1) below. By definition, GAG flows are those Gaussian flows whose ergodic self-joinings remain Gaussian. All Gaussian flows with simple spectrum are GAG flows [17]. If $\mathcal{T}^\sigma = (T_t^\sigma)_{t\in\mathbb{R}}$ denotes the Gaussian flow determined by a finite positive (continuous) measure σ on $\mathbb{R}_+$ and the flow is GAG then

$$I(\mathcal{T}^\sigma) \text{ is equal to the (multiplicative) group } -I(\sigma) \cup I(\sigma), \tag{1}$$

where $I(\sigma) = \{s \in \mathbb{R}_+^* : \sigma_s \equiv \sigma\}$ and $\sigma_s = (R_s)_*(\sigma)$ denotes the image of σ via the map $R_s : x \mapsto sx$ [17]. Recall that -1 is always a scale of self-similarity for a Gaussian flow.

In this note we focus on the problem of self-similarities in some subclasses of simple spectrum Gaussian flows. We first recall already known results. Classically, if σ is concentrated on an additively $\mathbb{Q}$-independent Borel set $A \subset \mathbb{R}_+$ then the Gaussian flow $\mathcal{T}^\sigma$ has simple spectrum, see [2]. Moreover, the subgroup $H := I(\mathcal{T}^\sigma) \cap \mathbb{R}_+^*$ is an additively $\mathbb{Q}$-independent set. Indeed, suppose that H is not an additively $\mathbb{Q}$-independent set. That is, for some distinct $h_1, \ldots, h_m \in H$ we have

$$\sum_{i=1}^{m} k_i h_i = 0 \quad \text{with} \quad k_i \in \mathbb{Z},\ i = 1, \ldots, m \quad \text{and} \quad \sum_{i=1}^{m} k_i^2 > 0. \tag{2}$$

Date: May 22, 2012.

[1] In [15] as well as in [17] only Gaussian automorphisms are considered, however all results can be rewritten for Gaussian flows.

Denote by $H_0 \subset H$ the multiplicative subgroup generated by $h_1, \ldots, h_m$. Since $H_0 \subset I(\mathcal{T}^\sigma)$, we have $\sigma_h \equiv \sigma$ for $h \in H_0$, thus the Borel set $B = \bigcap_{h \in H_0} hA$ has full σ-measure, is $\mathbb{Q}$-independent, and is literally H_0-invariant. Take any non-zero $x \in B$. Then the elements $h_i x \in B$, $i = 1, \ldots, m$, are distinct. Now, (2) yields

$$\sum_{i=1}^{m} k_i(h_i x) = x \sum_{i=1}^{m} k_i h_i = 0,$$

so B is not independent, a contradiction. On the other hand, in [6], it is shown that whenever a countable group $H \subset \mathbb{R}_+^*$ satisfies:

(3)
For each polynomial $P \in \mathbb{Q}[x_1, \ldots, x_m]$ if there is
a collection of distinct elements $h_1, \ldots, h_m$ in H such that
$P(h_1, \ldots, h_m) = 0$ then $P \equiv 0$,

then there exists a probability σ concentrated on a Borel $\mathbb{Q}$-independent set such that $I(\mathcal{T}^\sigma) = -H \cup H$. It is not difficult to see that the condition (3) is equivalent to saying that H is an additively $\mathbb{Q}$-independent set.

Theorem 1 ([6]). *Assume that $G = -H \cup H$, where $H \subset \mathbb{R}_+^*$ is a countable multiplicative subgroup. Then G can be realized as $I(\mathcal{T}^\sigma)$ for a Gaussian flow whose spectral measure σ is concentrated on a Borel $\mathbb{Q}$-independent set if and only if H is an additively $\mathbb{Q}$-independent set.*

Note that for H cyclic generated by $s \in \mathbb{R}_+$, the $\mathbb{Q}$-independence condition is equivalent to saying that s is transcendental. Hence, by Theorem 1, a real number s can be realized as a scale of self-similarity of a Gaussian flow whose spectral measure is concentrated on a $\mathbb{Q}$-independent Borel set if and only if s is transcendental.

On the other hand, there are no restrictions on H in the class of all Gaussian flows having simple spectrum.

Theorem 2 ([4]). *For each countable subgroup $H \subset \mathbb{R}_+^*$ there exists a simple spectrum Gaussian flow $\mathcal{T}^\sigma$ such that $I(\mathcal{T}^\sigma) = -H \cup H$.*

Note that, in particular, the above result of Danilenko and Ryzhikov brings the positive answer to the open problem [14] of existence of Gaussian flows $\mathcal{T}^\sigma$ with simple spectrum such that σ is not concentrated on a $\mathbb{Q}$-independent set; indeed, whenever H is not an additively $\mathbb{Q}$-independent set, by Theorem 1, the spectral measure σ resulting from Theorem 2 cannot be concentrated on a Borel $\mathbb{Q}$-independent set. See also [3] for constructions of Gaussian flows with zero entropy and having uncountable groups of self-similarities.

Our aim is to continue investigations on realization of countable subgroups as the groups of self-similarities in further restricted classes of Gaussian flows whose spectral measures are classical from the harmonic analysis point of view. Recall some basic notions. For every $s \in \mathbb{R}$ let $\xi_s : \mathbb{R} \to \mathbf{S}^1$ be given by $\xi_s(t) = \exp(2\pi i st)$. A finite positive Borel measure σ on $\mathbb{R}$ is called Kronecker if for each $f \in L^2(\mathbb{R}, \sigma)$, $|f| = 1$ σ-a.e., there exists a sequence $(t_n) \subset \mathbb{R}$, $t_n \to \infty$, such that

$$\xi_{t_n} \to f \quad \text{in} \quad L^2(\mathbb{R}, \sigma). \tag{4}$$

Each measure σ concentrated on a Kronecker set [12], [18] is a Kronecker measure. Indeed, Kronecker sets are compact subsets of $\mathbb{R}$ on which each continuous function of modulus one is a uniform limit of characters. Kronecker sets are examples of $\mathbb{Q}$-independent sets [18]. In general, as shown in [15], a Kronecker measure is concentrated on a Borel set which is the union of an increasing sequence of Kronecker sets, hence a Kronecker measure is concentrated on a Borel $\mathbb{Q}$-independent set, and the restriction on H in Theorem 1 applies. This turns out to be the only restriction as the main result of the note shows.

Theorem 3. *Assume that* $G = -H \cup H$, *where* $H \subset \mathbb{R}_+^*$ *is a countable multiplicative subgroup. Then* G *can be realized as* $I(\mathcal{T}^\sigma)$ [2] *for a Gaussian-Kronecker flow if and only if* H *is an additively* $\mathbb{Q}$*-independent set. In particular,* $h \in \mathbb{R}_+$ *can be a scale of self-similarity for a Gaussian-Kronecker flow if and only if* h *is transcendental.*

An extremal case when two dynamical systems are non-isomorphic is the disjointness in the Furstenberg sense [7], see also [9], [11], [14], [23] for disjointness results in ergodic theory. We would like also to emphasize that the notion of disjointness turned out to be quite meaningful in the problem of non-correlation with the Möbius function of sequences of dynamical origin [1]: we need that an automorphism T has the property that T^p and T^q are disjoint for any two different primes. In connection with that we will prove the following.

Theorem 4. *Assume that* $\mathcal{T}^\sigma = (T_t^\sigma)_{t\in\mathbb{R}}$ *is a Gaussian-Kronecker flow. If* $s \in \mathbb{Q}\setminus\{\pm 1\}$ *then* T_s^σ *is disjoint from* T_1^σ. *For every Gaussian-Kronecker automorphism* $T : (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ *the iterations* T^n, T^m *are disjoint for any two distinct natural numbers* n, m.

If s *is irrational then there exists a Gaussian-Kronecker flow* $\mathcal{T}^\sigma$ *such that* T_s^σ *and* T_1^σ *have a non-trivial common factor.*

An importance of Kronecker measures in ergodic theory follows from the following remarkable result of Foiaş and Stratila [5] (see also [2], and remarks on that result in [15] and [21]):

(5)
If $(S_t)_{t\in\mathbb{R}}$ is an ergodic flow of a standard probability
Borel space $(Y, \mathcal{C}, \nu)$, $f \in L^2(Y, \mathcal{C}, \nu)$ is real and the spectral measure
σ_f of f is the symmetrization of a Kronecker measure,
then the (stationary) process $(f \circ S_t)_{t\in\mathbb{R}}$ is Gaussian.

In [15], any measure σ satisfying the assertion (5) of Foiaş-Stratila theorem is called an FS measure. Each Kronecker measure is a Dirichlet measure[3] [18], but as shown in [15], there are FS measures which are not Dirichlet measures (see Figure 1). Moreover, in [21], it is announced that each continuous measure concentrated on an independent Helson[4] set is a Kronecker measure (for some examples in [21], the resulting Gaussian flows have no non-trivial rigid factors). We will strenghten Theorem 2 to the following result.



Figure 1. Different classes of measures

---

[2] In a sense, we can also control the flows $\mathcal{T}^{\sigma_s}$ for $s \notin -H \cup H$; we will prove their disjointness from $\mathcal{T}^\sigma$, see the proof of this theorem.

[3] A probability Borel measure σ on $\mathbb{R}$ is Dirichlet, if (4) is satisfied for $f = 1$. From the dynamical point of view, Dirichlet measures correspond to rigidity: a flow $\mathcal{T}$ is *rigid* if $T^{t_n} \to Id$ for some $t_n \to \infty$.

[4] $A \subset \mathbb{R}$ is called Helson if for some $\delta > 0$ and each complex Borel measure κ concentrated on A the $\sup_{t\in\mathbb{R}} \left|\int_{\mathbb{R}} e^{2\pi itx}\, d\kappa(x)\right|$ is bounded away from the δ-fraction of the total variation of κ.

Theorem 5. *Any symmetric countable group* $G \subset \mathbb{R}^*$ *can be realized as the group of self-similarities of a simple spectrum Gaussian flow* $\mathcal{T}^\sigma$ *with* σ *being an FS measure.*

In particular, in connection with the forementioned question from [14], there is an FS measure for which the Gaussian flow has simple spectrum but σ is not concentrated on a $\mathbb{Q}$-independent set. These are apparently the first examples of FS measures which are not concentrated on $\mathbb{Q}$-independent Borel sets but yield Gaussian flows with simple spectrum (cf. [15] and [21]).

At the end of the note we will discuss self-similarity properties of Gaussian flows arising from a "typical" measure or from the maximal spectral types of a "typical" flow (cf. the disjointness results from [4]).

Theorem 6. *Assume that* $0 \leq a < b$. *For a "typical"* $\sigma \in \mathcal{P}([a,b])$ *the flow* $\mathcal{T}^\sigma$ *is Gaussian-Kronecker such that for each* $|r| \neq |s|$ *the flows* $\mathcal{T}^{\sigma_r}$ *and* $\mathcal{T}^{\sigma_s}$ *are disjoint. In particular* $I(\mathcal{T}^\sigma) = \{\pm 1\}$.

For a "typical" flow $\mathcal{T}$ *of a standard probability Borel space* $(X, \mathcal{B}, \mu)$, *for its maximal spectral type* $\sigma_{\mathcal{T}}$ *we have:* $\mathcal{T}^{\sigma_{\mathcal{T}}|_{\mathbb{R}_+}}$ *has simple spectrum and for* $|r| \neq |s|$ *the flows* $\mathcal{T}^{(\sigma_{\mathcal{T}}|_{\mathbb{R}_+})_r}$ *and* $\mathcal{T}^{(\sigma_{\mathcal{T}}|_{\mathbb{R}_+})_s}$ *are disjoint. In particular* $I(\mathcal{T}^{\sigma_{\mathcal{T}}|_{\mathbb{R}_+}}) = \{\pm 1\}$.

1. NOTATION AND BASIC RESULTS

Assume that $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ is a measurable[5] measure-preserving flow acting on a standard probability Borel space $(X, \mathcal{B}, \mu)$. It then induces a (continuous) one-parameter group of unitary operators acting on $L^2(X, \mathcal{B}, \mu)$ by the formula $T_t f = f \circ T_t$. By Bochner's theorem, the function $t \mapsto \int_X T_t f \cdot \overline{f}\, d\mu$ determines the so called *spectral measure* σ_f of f for which $\widehat{\sigma}_f(t) = \int_X T_t f \cdot \overline{f}\, d\mu$, $t \in \mathbb{R}$. Usually, one only considers spectral measures of $f \in L^2_0(X, \mathcal{B}, \mu)$, that is, of elements with zero mean (the spectral measure of the constant function c is equal to $|c|^2\delta_0$). Then σ_f is a finite positive Borel measure on $\mathbb{R}$. Among spectral measures there are maximal ones with respect to the absolute continuity relation. Each such maximal measure is called a *maximal spectral type measure* and, by some abuse of notation, it will be denoted by $\sigma_{\mathcal{T}}$. We refer the reader to [11] and [14] for some basics about spectral theory of unitary representations of locally compact Abelian groups in the dynamical context.

Assume that $\mathcal{T}$ is ergodic and let $\mathcal{S} = (S_t)_{t\in\mathbb{R}}$ be another ergodic flow (acting on $(Y, \mathcal{C}, \nu)$). Any probability measure ρ on $(X \times Y, \mathcal{B} \otimes \mathcal{C})$ which is $(T_t \times S_t)_{t\in\mathbb{R}}$-invariant and has marginals μ and ν respectively, is called a *joining* of $\mathcal{T}$ and $\mathcal{S}$. If, additionally, the flow $((T_t \times S_t)_{t\in\mathbb{R}}, \rho)$ is ergodic then ρ is called an *ergodic joining* [6]. The ergodic joinings are extremal points in the simplex of all joinings. If the set of joinings is reduced to contain only the product measure then one speaks about *disjointness* of $\mathcal{T}$ and $\mathcal{S}$ [7] and we will write $\mathcal{T} \perp \mathcal{S}$. Similar notions appear when one considers automorphisms. Note that whenever for some $t \neq 0$, $T_t \perp S_t$ then $\mathcal{T} \perp \mathcal{S}$. Note also that whenever

$$\mathcal{T} \text{ is weakly mixing then } \mathcal{T} \perp \mathcal{S} \text{ if and only if } T_1 \perp S_1. \tag{6}$$

Indeed, if $T_1 \not\perp S_1$ then there exists an ergodic joining ρ between them different than the product measure. Then, $\rho \circ (T_r \times S_r)$ for $0 \leq r < 1$ has the same properties. By disjointness of $\mathcal{T}$ and $\mathcal{S}$, $\int_0^1 \rho \circ (T_r \times S_r)\, dr = \mu \otimes \nu$. But T_1 is weakly mixing, so $\mu \otimes \nu$ is an ergodic joining of T_1 and S_1, and therefore $\rho \circ (T_r \times S_r) = \mu \otimes \nu$. We refer the reader to [9] for the theory of joinings in ergodic theory.

A flow $\mathcal{T}$ is called Gaussian if there is a $\mathcal{T}$-invariant subspace $\mathcal{H} \subset L^2_0(X, \mathcal{B}, \mu)$ of the zero mean real-valued functions such that all non-zero variables from $\mathcal{H}$ are

---

[5] Measurability means that for each $f \in L^2(X, \mathcal{B}, \mu)$ the map $t \mapsto f \circ T_t$ is continuous.

[6] If $\mathcal{T} = \mathcal{S}$ then we speak about *self-joinings*.

Gaussian and the smallest σ-algebra making all these variables measurable equals $\mathcal{B}$. A Gaussian flow is ergodic if and only if the maximal spectral type on $\mathcal{H}$ is continuous (and then $\mathcal{T}$ is weakly mixing). Since Gaussian variables are real, it is not hard to see that their spectral measures are symmetric, that is, for $f \in \mathcal{H}$, σ_f is invariant under the map $R_{-1} : x \mapsto -x$.

A standard way to obtain a (weakly mixing) Gaussian flow is to start with a finite positive continuous Borel measure σ on $\mathbb{R}_+$. Consider the symmetrization $\widetilde{\sigma} = \sigma + (R_{-1})_*\sigma$ [7]. We let $\mathcal{V} = (V_t)_{t\in\mathbb{R}}$ denote the one-parameter group of unitary operators on $L^2(\mathbb{R}, \widetilde{\sigma})$ defined by $V_t(f)(x) = e^{2\pi itx}f(x)$. Then the correspondence

$$f(x) \mapsto f(-x) \tag{7}$$

yields the unitary conjugation of $\mathcal{V}$ and its inverse. Let $(X, \mathcal{B}, \mu)$ be a *Gaussian probability space*, that is, a standard probability space together with an infinite dimensional, closed, real and $\mathcal{B}$-generating subspace $\mathcal{H} \subset L^2(X, \mathcal{B}, \mu)$ whose all non-zero variables are Gaussian. We then consider $\mathcal{H} + i\mathcal{H}$, so called *complex Gaussian space*, and define an isomorphic copy of $\mathcal{V}$ on it. It is then standard to show (see e.g. [17], Section 2) that $\mathcal{V}$ has a unique extension to a (measurable) flow $\mathcal{T}^\sigma = (T^\sigma_t)$ of $(X, \mathcal{B}, \mu)$ for which $U_{T^\sigma_t}|_{\mathcal{H}+i\mathcal{H}} = V_t$, $t \in \mathbb{R}$. By the same token, the correspondence (7) extends to an isomorphism of $(X, \mathcal{B}, \mu)$ which conjugates the Gaussian flow and its inverse $(T^\sigma_{-t})_{t\in\mathbb{R}}$.

A Gaussian flow $\mathcal{T}^\sigma$ is called Gaussian-Kronecker (FS resp.) if σ is a continuous Kronecker (FS resp.) measure. Following [17], a Gaussian flow $\mathcal{T}^\sigma$ (with the Gaussian space $\mathcal{H}$) is called GAG if for each its ergodic self-joining η the space

$$\{f(x) + g(y) : f, g \in \mathcal{H}\}$$

consists solely of Gaussian variables (the flow $(T^\sigma_t \times T^\sigma_t)_{t\in\mathbb{R}}$ is then a Gaussian flow as well). We have [17]



FIGURE 2. Different subclasses of GAG flows

For all these classes of flows we have that if $\mathcal{T}^\sigma$ is in the class, so is $\mathcal{T}^{\sigma_s}$ for $s \neq 0$.

In general, Gaussian flows given by equivalent measures are isomorphic. It follows from [17] that any isomorphism between a GAG flow $\mathcal{T}^\sigma$ and another Gaussian flow $\mathcal{T}^\nu$ is entirely determined by a unitary isomorphism of restrictions of the unitary actions $(T^\sigma_t)_{t\in\mathbb{R}}$ and $(T^\nu_t)_{t\in\mathbb{R}}$ to their Gaussian subspaces. That is, in the GAG situation, $\mathcal{T}^\sigma$ are $\mathcal{T}^\nu$ are isomorphic if and only if $\sigma \equiv \nu$. If we apply that to σ and σ_s for $s \in \mathbb{R}_+$ we will immediately get (1) to hold (in the GAG case).

We will now prove the following.

Proposition 7. *Assume that $\mathcal{T}^\sigma$ is GAG. Fix $s \neq 0$. Then the sets of self-joinings of $\mathcal{T}^\sigma$ and of self-joinings of T^σ_s are the same. (Hence ergodic self-joinings are also the same.) In particular, the factors and the centralizer of the flow and of the time s-automorphism are the same.*

---

[7] In general, when f is a measurable map from $(X, \mathcal{B})$ to $(Y, \mathcal{C})$ and κ is a probability measure on X then $f_*(\kappa)$ is the measure on Y defined by $f_*(\kappa)(C) = \kappa(f^{-1}(C))$.

Proof. This follows from the proof of Theorem 6.1 in [10] which asserts that such an equality of the sets of self-joinings takes place whenever each ergodic self-joining of the flow is an ergodic self-joining for the time-s automorphism. In the GAG case, by definition, such ergodic joinings for the flow $\mathcal{T}^\sigma$ are Gaussian joinings, so they are automatically ergodic for the T^σ_s [17]. □

Corollary 8. *Assume that $\mathcal{T}^\sigma$ is GAG. Then T^σ_s is a GAG automorphism for each $s \neq 0$.*

We will also make use of the following results.

Theorem 9 ([17])**.** *Assume that $\mathcal{T}^\sigma$ is GAG and let $\mathcal{T}^\eta$ be an arbitrary Gaussian flow. Then $\mathcal{T}^\sigma \perp \mathcal{T}^\eta$ if and only if $\widetilde{\sigma} \perp \widetilde{\eta} * \delta_r$ for each $r \in \mathbb{R}$.*

Proposition 10 ([15])**.** *If σ_1 and σ_2 are measures with the FS property and $\mathcal{T}^{\sigma_1} \perp \mathcal{T}^{\sigma_2}$ then $\sigma = \frac{1}{2}(\sigma_1 + \sigma_2)$ is also an FS measure.*

2. Auxiliary lemmas

Given a compact subset $X \subset \mathbb{R}$ denote by $\mathcal{P}(X)$ the set of all Borel probability measures concentrated on X endowed with the usual weak topology which is compact and metrizable: if $\{f_n : n \geq 1\}$ is a dense set in $C(X)$ then

$$d(\sigma,\eta) = \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{\left|\int f_n\,d\sigma - \int f_n\,d\eta\right|}{1 + \left|\int f_n\,d\sigma - \int f_n\,d\eta\right|} \tag{8}$$

defines a metric compatible with the weak topology. Denote $\mathcal{U}(X) = \{f \in C(X) : |f| = 1\}$ which is a closed subset of $C(X)$ in the uniform topology, in particular $\mathcal{U}(X)$ is a Polish space.

Lemma 11. *Assume that $X = [a,b]$. Let $\{h_0, h_1, \ldots, h_m\} \subset \mathbb{R}^*$ be a $\mathbb{Q}$-independent set. Then for each $f \in \mathcal{U}\left(\bigcup_{j=0}^m h_j X\right)$ and $\varepsilon > 0$*

$$\begin{gathered} A_{f,\varepsilon}(h_1,\ldots,h_m) := \\ \left\{\sigma \in \mathcal{P}([a,b]) : (\exists t \in \mathbb{R})\ \|f - \xi_t\|_{L^2\left(\mathbb{R}, \frac{1}{m+1}\sum_{j=0}^m \sigma_{h_j}\right)} < \varepsilon\right\} \end{gathered} \tag{9}$$

is open and dense in $\mathcal{P}([a,b])$.

Proof. The set $A_{f,\varepsilon}(h_1,\ldots,h_m)$ is clearly open, so we need to show its denseness in $\mathcal{P}(X)$. Since discrete measures with a finite number of atoms form a dense subset of $\mathcal{P}(X)$ we take $\nu = \sum_{s=1}^N a_s \delta_{y_s}$ with $y_s \in [a,b]$, $a_s > 0$, $s = 1,\ldots,N$ and $\sum_{s=1}^N a_s = 1$ and fix $\delta > 0$. All we need to show is to find a subset $\{x_1,\ldots,x_N\} \subset [a,b]$ such that $|x_s - y_s| < \delta$ for $s = 1,\ldots,N$ and such that the set

$$L := \bigcup_{j=0}^m \{h_j x_1, \ldots, h_j x_N\} \quad \text{is } \mathbb{Q}\text{-independent.}$$

Indeed, in this case by Kronecker's theorem, the set L is a finite Kronecker set, so the measure $\frac{1}{m+1}\sum_{j=0}^m \left(\sum_{s=1}^N a_s \delta_{x_s}\right)_{h_j}$ is Kronecker, whence belongs to $A_{f,\varepsilon}(h_1,\ldots,h_m)$ and it δ-approximates ν. To show that $x_1,\ldots,x_N$ can be selected so that L is $\mathbb{Q}$-independent, consider the algebraic varieties of the form

$$\left\{(z_1,\ldots,z_N) \in X^{\times N} : \sum_{j=0}^m \sum_{s=1}^N k_{js} h_j z_s = 0\right\}$$

for some non-zero integer matrix (k_{js}). Since

$$\sum_{j=0}^{m}\sum_{s=1}^{N} k_{js}h_j z_s = \sum_{s=1}^{N}\left(\sum_{j=0}^{m} k_{js}h_j\right) z_s$$

and $\sum_{j=0}^{m} k_{js}h_j \neq 0$ whenever $(k_{0s},\ldots,k_{ms}) \neq (0,\ldots,0)$ (and there are such vectors since the matrix (k_{js}) is not zero), each such variety has N-dimensional Lebesgue measure zero. Since there are only countably many such varieties involved, we may discard the union S of them from $[a,b]^{\times N}$. Now, each choice of $(x_1,\ldots,x_N)$ from $(y_1-\delta,y_1+\delta)\times\ldots\times(y_N-\delta,y_N+\delta)\setminus S$ satisfies our requirements. $\square$

Lemma 12. *Given $H \subset \mathbb{R}_+^*$ a countable subset which is a $\mathbb{Q}$-independent set, the set of continuous (Kronecker) measures $\sigma \in \mathcal{P}([a,b])$ for which the measure*

$$\sum_{h\in H} a_h\sigma_h \quad \textit{is a Kronecker measure (on } \mathbb{R}) \tag{10}$$

for each choice of $a_h \geq 0$, $\sum_{h\in H} a_h = 1$, is a G_δ and dense subset of $\mathcal{P}([a,b])$.

Proof. Denote by $\mathcal{P}_c([a,b])$ the set of continuous measures which is a G_δ and dense subset of $\mathcal{P}([a,b])$. Let $H=\{h_0,h_1,h_2,\ldots\}$. For every $m\geq 0$ fix a countable dense family $\left\{g_i^{(m)} : i\geq 1\right\} \subset \mathcal{U}\left(\bigcup_{i=0}^{m} h_i[a,b]\right)$. Then, by Lemma 11, the set

$$\mathcal{P}_c([a,b]) \cap \bigcap_{m=1}^{\infty}\bigcap_{i=1}^{\infty}\bigcap_{p=1}^{\infty} A_{g_i^{(m)},\frac{1}{p}}(h_1,\ldots,h_m)$$

is G_δ and dense in $\mathcal{P}([a,b])$ and it remains to show that this is precisely the set of measures satisfying (10). Indeed, given $m\geq 1$, the set

$$\mathcal{K}_m(H) := \mathcal{P}_c([a,b]) \cap \bigcap_{i=1}^{\infty}\bigcap_{p=1}^{\infty} A_{g_i^{(m)},\frac{1}{p}}(h_1,\ldots,h_m)$$

is precisely the set of continuous Kronecker measures $\sigma \in \mathcal{P}([a,b])$ such that the measure $\frac{1}{m+1}\sum_{i=0}^{m}\sigma_{h_i}$ is a Kronecker measure (on the real line). Moreover, each measure absolutely continuous with respect to a Kronecker measure is also a Kronecker measure [15]. Therefore the set $\mathcal{K}_m(H)$ is equal to the set of all Kronecker measures $\sigma\in\mathcal{P}([a,b])$ such that $\sum_{i=0}^{m} b_i\sigma_{h_i}$ is Kronecker for arbitrary choice of $b_i\geq 0$, $\sum_{i=0}^{m} b_i = 1$. Finally, for each $m\geq 1$,

$$\frac{1}{\sum_{i=0}^{m} a_{h_i}}\sum_{i=0}^{m} a_{h_i}\sigma_{h_i} \ll \frac{1}{m+1}\sum_{i=0}^{m}\sigma_{h_i},$$

so if for each $m\geq 1$ the measure $\frac{1}{m+1}\sum_{i=0}^{m}\sigma_{h_i}$ is Kronecker, so is $\sum_{h\in H} a_h\sigma_h$. $\square$

Remark 1. The idea of the above proofs is taken from a letter that has been sent to us by T.W. Körner. In this letter, T.W. Körner shows that given a transcendental number $h\in\mathbb{R}$, for a "typical" (in the Hausdorff metric) closed subset $K\subset[a,b]$ the set $K\cup hK$ is Kronecker and uncountable. The proofs are the same since finite sets are dense in the Hausdorff metric and if h is transcendental then given distinct $y_1,\ldots,y_N\in[a,b]$ and $\delta>0$ we can find $q_i\in\mathbb{Q}$ so that for $x_i := h^{2i}q_i$ we have $|x_i-y_i|<\delta$ for $i=1,\ldots,N$ and clearly the set $\{x_1,\ldots,x_N,hx_1,\ldots,hx_N\}$ is $\mathbb{Q}$-independent. It only remains to notice that uncountable closed subsets are typical in the Hausdorff metric.

Note also that using the proofs of Lemmas 11 and 12, given $H\subset\mathbb{R}_+^*$ a countable multiplicative subgroup which is additively $\mathbb{Q}$-independent, we obtain that a typical (with respect to the Hausdorff distance) closed subset $K\subset[a,b]$ has the property that for each finite subset $C \subset H$ the set $\bigcup_{h\in C} hK$ is Kronecker, so the set $\bigcup_{h\in H} hK$ is a $\mathbb{Q}$-independent F_σ-set.

We will also need the following "compact $\mathbb{Q}$-independent set" version of Lemma 12.

Lemma 13. *Assume that $K \subset \mathbb{R}$ is a compact uncountable set. Assume that $H \subset \mathbb{R}_+^*$ is a countable set which is additively $\mathbb{Q}$-independent. Assume moreover that the set $\bigcup_{h\in H} hK$ is $\mathbb{Q}$-independent. Then the set of continuous (Kronecker) measures σ concentrated on K for which the measure*

$$\sum_{h\in H} a_h\sigma_h \quad \textit{is a Kronecker measure} \tag{11}$$

for each choice of $a_h \geq 0$, $\sum_{h\in H} a_h = 1$, is a G_δ and dense subset of $\mathcal{P}(K)$.

Proof. This follows from the proofs of Lemmas 11 and 12, where in addition the proof of Lemma 11 is simplified; indeed, for any choice of $\{y_1, \ldots, y_N\} \subset K$ the set $\bigcup_{j=0}^{m}\{h_j y_1, \ldots, h_j y_N\}$ is $\mathbb{Q}$-independent by assumption (so we may take $x_i = y_i$ for $i = 1, \ldots, N$). □

Remark 2. For any non-trivial compact interval $[a, b] \subset \mathbb{R}$ denote by $\mathcal{P}_c^{[a,b]}(\mathbb{R})$ the set of measures $\nu \in \mathcal{P}_c(\mathbb{R})$ such that $\nu([a, b]) > 0$. Since the map $\mathcal{P}_c(\mathbb{R}) \ni \nu \mapsto \nu([a, b]) \in \mathbb{R}$ is continuous, the set $\mathcal{P}_c^{[a,b]}(\mathbb{R})$ is open and dense in $\mathcal{P}_c(\mathbb{R})$. Let us consider the map $\Delta = \Delta^{[a,b]} : \mathcal{P}_c^{[a,b]}(\mathbb{R}) \to \mathcal{P}_c([a, b])$ such that $\Delta(\nu)$ is the conditional probability measure $\nu(\,\cdot\,|[a, b])$. This map is continuous and the preimage of any dense subset of $\mathcal{P}_c([a, b])$ is dense in $\mathcal{P}_c^{[a,b]}(\mathbb{R})$. Indeed, let $A \subset \mathcal{P}_c([a, b])$ be dense and take any $\nu \in \mathcal{P}_c^{[a,b]}(\mathbb{R})$. Then there exists a sequence $(\widetilde{\nu}_n)_{n\leq 1}$ in A such that $\widetilde{\nu}_n \to \Delta(\nu)$ weakly. For every $n \geq 1$ define $\nu_n \in \mathcal{P}_c^{[a,b]}(\mathbb{R})$ so that the restriction of ν_n to $[a, b]$ is $\nu([a, b])\widetilde{\nu}_n$ and the measures ν_n and ν coincide on $\mathbb{R} \setminus [a, b]$. Then $\Delta(\nu_n) = \widetilde{\nu}_n \in A$ and $\nu_n \to \nu$ weakly. Consequently, the preimage $\Delta^{-1}A$ of any G_δ dense subset $A \subset \mathcal{P}_c([a, b])$ is G_δ dense in $\mathcal{P}_c^{[a,b]}(\mathbb{R})$.

Before we prove a certain disjointness property of Kronecker measures, we will need the following general observation.

Lemma 14. *Let $(X, \mathcal{B})$ be a standard Borel space and let $\varphi : X \to X$ be a measurable map. Let σ be a finite positive continuous Borel measure on X such that the map $\varphi : (X, \sigma) \to (X, \varphi_*\sigma)$ is almost everywhere invertible. Assume that $\sigma(\{x \in X : \varphi(x) = x\}) = 0$ and that the measures σ and $\varphi_*\sigma$ are not mutually singular. Then there exists a measurable set $A \in \mathcal{B}$ such that $\sigma(A) > 0$, $\sigma(A \cap \varphi^{-1}A) = 0$ and the measures σ and $\varphi_*\sigma$ restricted to A are equivalent.*

Proof. By assumption, there exists $Y \in \mathcal{B}$ such that $\sigma(Y) > 0$ and the measures σ and $\varphi_*\sigma$ restricted to Y are equivalent. It follows that if $A \in \mathcal{B}$, $A \subset Y$, $\sigma(A) > 0$, then the measures σ and $\varphi_*\sigma$ restricted to A are also equivalent.

Case 1. Suppose that there exists $B \in \mathcal{B}$ such that $B \subset Y$ and $\sigma(B \setminus \varphi(B)) > 0$. Set $A := B \setminus \varphi(B)$. Then $\sigma(A) > 0$ and $A \cap \varphi^{-1}A = (B \setminus \varphi(B)) \cap (\varphi^{-1}B \setminus B) = \emptyset$. Since $A \subset B \subset Y$, our claim follows.

Case 2. Suppose that for every $B \in \mathcal{B}$ with $B \subset Y$ we have $\sigma(B \setminus \varphi(B)) = 0$. As σ and $\varphi_*\sigma$ restricted to Y are equivalent, it follows that

$$0 = \varphi_*\sigma(B \setminus \varphi(B)) = \sigma(\varphi^{-1}B \setminus B) \quad \text{for every} \quad B \subset Y. \tag{12}$$

We now show that there exists $A \in \mathcal{B}$ such that $A \subset Y$, $\sigma(A) > 0$ and $\sigma(A\cap\varphi^{-1}A) = 0$, which gives our assertion. Suppose that, contrary to our claim, for every $A \in \mathcal{B}$ with $A \subset Y$ the condition $\sigma(A) > 0$ implies $\sigma(A \cap \varphi^{-1}A) > 0$. It follows that

$$\sigma(B \setminus \varphi^{-1}B) = 0 \quad \text{for every measurable} \quad B \in \mathcal{B} \text{ with } B \subset Y. \tag{13}$$

Indeed, otherwise for some B as above, $A := B \setminus \varphi^{-1}(B) \subset Y$ would be of positive σ-measure and since

$$(B \setminus \varphi^{-1}B) \cap \varphi^{-1}(B \setminus \varphi^{-1}B) = (B \setminus \varphi^{-1}B) \cap (\varphi^{-1}B \setminus \varphi^{-2}B) = \emptyset,$$

and we would get $\sigma(A \cap \varphi^{-1}A) = 0$, a contradiction.

Now, (12) combined with (13) gives $\sigma(B \triangle \varphi^{-1}B) = 0$ for every $B \in \mathcal{B}$ with $B \subset Y$. It follows that $\varphi(x) = x$ for σ-a.e. $x \in Y$, contrary to assumption. □

For any real s let $\theta_s : \mathbb{R} \to \mathbb{R}$, $\theta_s(t) = t + s$. Recall that for every $n \in \mathbb{Z}$ and $z_1, z_2 \in \mathbf{S}^1$ we have

$$|z_1^n - z_2^n| \leq |n||z_1 - z_2|. \tag{14}$$

Lemma 15. *Let σ be a continuous Kronecker measure on $\mathbb{R}$. Then for every $s \in \mathbb{Q}^* \setminus \{1\}$ and $r \in \mathbb{R}$ we have $\sigma \perp \sigma_s * \delta_r$.*

Proof. Suppose that, contrary to our claim, there exists $s \in \mathbb{Q}^* \setminus \{1\}$ and $r \in \mathbb{R}$ such that $\sigma \not\perp \sigma_s * \delta_r$. Let $\varphi := \theta_r \circ R_s$. Then $\varphi : \mathbb{R} \to \mathbb{R}$ is an invertible map with one fixed point and $\sigma_s * \delta_r = \varphi_* \sigma$. By Lemma 14, there exists a Borel set $A_0 \subset \mathbb{R}$ such that $\sigma(A_0) > 0$, $\sigma(A_0 \cap \varphi^{-1}A_0) = 0$ and the measures σ, $\varphi_*\sigma$ restricted to A_0 are equivalent. Thus $\sigma(\varphi^{-1}A_0) > 0$. Let $A_1, A_2 \subset A_0$ be disjoint Borel subsets such that $\sigma(\varphi^{-1}A_1) > 0$ and $\sigma(\varphi^{-1}A_2) > 0$.

Let $s = q/p$ with p and q relatively prime integer numbers. Choose $z_0 \in \mathbf{S}^1$ such that $z_0^q \neq 1$. Let us consider the measurable map $f : \mathbb{R} \to \mathbf{S}^1$ such that $f(x) = z_0$ if $x \in \varphi^{-1}A_2$ and $f(x) = 1$ otherwise. Since σ is a Kronecker measure, there exists a sequence $(t_n)_{n \in \mathbb{N}}$ of real numbers such that $\xi_{t_n} \to f$ in $L^2(\mathbb{R}, \sigma)$. Thus $\xi_{t_n} \circ \varphi^{-1} \to f \circ \varphi^{-1}$ in $L^2(\mathbb{R}, \varphi_*\sigma)$. Since

$$\begin{aligned}
g_n^0(x) &:= \chi_{A_0}(x) \left|\exp(2\pi i t_n x) - 1\right| \leq |\xi_{t_n}(x) - f(x)| \\
g_n^1(x) &:= \chi_{A_1}(x) \left|\exp(2\pi i t_n s^{-1}(x - r)) - 1\right| \leq \left|\xi_{t_n}(\varphi^{-1}x) - f(\varphi^{-1}x)\right| \\
g_n^2(x) &:= \chi_{A_2}(x) \left|\exp(2\pi i t_n s^{-1}(x - r)) - z_0\right| \leq \left|\xi_{t_n}(\varphi^{-1}x) - f(\varphi^{-1}x)\right|,
\end{aligned}$$

it follows that (g_n^0) tends to zero in measure σ and the sequences (g_n^1), (g_n^2) tend to zero in measure $\varphi_*\sigma$. As $\sigma \equiv \varphi_*\sigma$ on A_0 and $A_1, A_2 \subset A_0$, the sequences (g_n^1), (g_n^2) tend to zero in measure σ, as well. Fix

$$0 < \varepsilon < \frac{|z_0^q - 1|}{2(|p| + |q|)}. \tag{15}$$

Then there exist measurable sets $A_k' \subset A_k$, $k = 0, 1, 2$ and $n \in \mathbb{N}$ such that for $k = 0, 1, 2$

$$\sigma(A_k \setminus A_k') < \frac{1}{4} \min(\sigma(A_1), \sigma(A_2)) \text{ and } g_n^k(x) < \varepsilon \text{ for all } x \in A_k'.$$

Therefore for $k = 1, 2$ we have

$$\sigma(A_k \setminus A_0') \leq \sigma(A_0 \setminus A_0') < \frac{1}{4}\sigma(A_k) \text{ and } \sigma(A_k \setminus A_k') < \frac{1}{4}\sigma(A_k),$$

so $\sigma(A_0' \cap A_k') > \sigma(A_k)/2 > 0$. Choose two real numbers $x_1 \in A_0' \cap A_1'$ and $x_2 \in A_0' \cap A_2'$. Then

$$\begin{aligned}
&|\exp(2\pi i t_n x_1) - 1| = g_n^0(x_1) < \varepsilon, \quad \left|\exp(2\pi i t_n \frac{p}{q}(x_1 - r)) - 1\right| = g_n^1(x_1) < \varepsilon, \\
&|\exp(2\pi i t_n x_2) - 1| = g_n^0(x_2) < \varepsilon, \quad \left|\exp(2\pi i t_n \frac{p}{q}(x_2 - r)) - z_0\right| = g_n^2(x_2) < \varepsilon.
\end{aligned}$$

In view of (14),

$$\begin{aligned} &|\exp(2\pi i t_n p\, x_1) - 1| < |p|\varepsilon, \quad |\exp(2\pi i t_n p(x_1 - r)) - 1| < |q|\varepsilon, \\ &|\exp(2\pi i t_n p\, x_2) - 1| < |p|\varepsilon, \quad |\exp(2\pi i t_n p(x_2 - r)) - z_0^q| < |q|\varepsilon. \end{aligned}$$

Hence

$$|\exp(2\pi i t_n pr) - 1| < (|p| + |q|)\varepsilon, \quad |\exp(2\pi i t_n pr) z_0^q - 1| < (|p| + |q|)\varepsilon,$$

so

$$|1 - z_0^q| < 2(|p| + |q|)\varepsilon,$$

contrary to (15). □

Let us now consider the space $\mathcal{P}(\mathbb{R})$ of all Borel probability measures on $\mathbb{R}$ endowed with the weak topology.

By $\mathrm{supp}(\sigma)$ we always mean the topological support of the measure σ. Let us recall that

$$\begin{aligned} &\text{if } \sigma \in \mathcal{P}(\mathbb{R}) \text{ has } \mathrm{supp}(\sigma) = \mathbb{R} \\ &\text{then the set } \{\nu \in \mathcal{P}(\mathbb{R}) \colon \nu \ll \sigma\} \text{ is dense in } \mathcal{P}(\mathbb{R}). \end{aligned} \tag{16}$$

Denote by $\mathcal{P}_c(\mathbb{R})$ the set of all continuous members of $\mathcal{P}(\mathbb{R})$ (this is a G_δ and dense subset of $\mathcal{P}(\mathbb{R})$).

The proof of the lemma below is a slight modification of the proof of Lemma 3.1 from [4].

Lemma 16. *The set*

$$\mathcal{S} = \{\sigma \in \mathcal{P}_c(\mathbb{R}) \colon \sigma_s \perp \sigma * \delta_t \quad \textit{for each}\, 1 \neq s \in \mathbb{R}^*,\ t \in \mathbb{R}\}$$

is G_δ and dense in $\mathcal{P}(\mathbb{R})$.

Proof. Denote by $\mathcal{I}$ the family of open subset of $\mathbb{R}$ which are finite unions of open intervals. Recall that for two measures $\sigma, \nu \in \mathcal{P}(\mathbb{R})$

$$\sigma \perp \nu \iff \forall_{n\in\mathbb{N}}\ \exists_{\mathcal{O}\in\mathcal{I}}\ \sigma(\mathcal{O}) < 1/n \quad \text{and} \quad \nu(\mathcal{O}) > 1 - 1/n. \tag{17}$$

For any compact rectangle $I \times J \subset (\mathbb{R}^* \setminus \{1\}) \times \mathbb{R}$ denote by $\mathcal{V}(I \times J)$ the set of all finite covers of $I \times J$ by compact rectangles contained in $(\mathbb{R}^* \setminus \{1\}) \times \mathbb{R}$. Notice that for each open subset $\mathcal{O} \in \mathcal{I}$ the map

$$\mathcal{P}_c(\mathbb{R}) \times \mathbb{R}^* \times \mathbb{R} \ni (\sigma, s, r) \mapsto \sigma_s * \delta_r(\mathcal{O}) \in \mathbb{R} \tag{18}$$

is continuous. Therefore, given a compact rectangle $F \subset (\mathbb{R}^* \setminus \{1\}) \times \mathbb{R}$ and an open subset $\mathcal{O} \in \mathcal{I}$ the map

$$f_{F,\mathcal{O}} \colon \mathcal{P}_c(\mathbb{R}) \ni \sigma \mapsto \Big(\sigma(\mathcal{O}), \max_{(s,r)\in F} \sigma_s * \delta_r(\mathcal{O})\Big) \in \mathbb{R}^2$$

is continuous. Let

$$\widetilde{\mathcal{S}} = \bigcap_{I \not\ni 1} \bigcap_{J} \bigcap_{n\in\mathbb{N}} \bigcup_{\kappa\in\mathcal{V}(I\times J)} \bigcap_{F\in\kappa} \bigcup_{\mathcal{O}\in\mathcal{I}} f_{F,\mathcal{O}}^{-1}\left((1 - 1/n, \infty) \times (-\infty, 1/n)\right),$$

where I and J run over closed intervals with rational endpoints. Then $\widetilde{\mathcal{S}}$ is a G_δ set.

We claim that $\widetilde{\mathcal{S}} = \mathcal{S}$. Indeed, let $\sigma \in \mathcal{S}$. Let $I \not\ni 1$ and $J \subset \mathbb{R}$ be compact intervals and $n \in \mathbb{N}$. By assumption and (17), for every $(s_0, r_0) \in I \times J$ there exists an open set $\mathcal{O}_{s_0,r_0} \in \mathcal{I}$ such that

$$\sigma(\mathcal{O}_{s_0,r_0}) > 1 - 1/n \quad \text{and} \quad \sigma_{s_0} * \delta_{r_0}(\mathcal{O}_{s_0,r_0}) < 1/n.$$

Since the map (18) is continuous, there exist open rectangles $U'_{s_0,r_0} \subset U_{s_0,r_0} \subset \mathbb{R}^2$ such that $(s_0, r_0) \in U'_{s_0,r_0}$ and a compact rectangle $F_{s_0,r_0} \subset (\mathbb{R}^* \setminus \{1\}) \times \mathbb{R}$ satisfying $U'_{s_0,r_0} \subset F_{s_0,r_0} \subset U_{s_0,r_0}$ such that

$$\sigma_s * \delta_r(\mathcal{O}_{s_0,r_0}) < 1/n \quad \text{for all } (s,r) \in U_{s_0,r_0}.$$

Since $I \times J$ is compact and $\{U'_{s,r} : (s,r) \in I \times J\}$ is its open cover, there exists a finite cover $\kappa := \{F_{s_1,r_1}, \ldots, F_{s_k,r_k}\}$ of $I \times J$. It follows that

$$f_{F_{s_j,r_j}, \mathcal{O}_{s_j,r_j}}(\sigma) \in (1 - 1/n, \infty) \times (-\infty, 1/n) \quad \text{for all} \quad j = 1, \ldots, k,$$

thus $\sigma \in \widetilde{\mathcal{S}}$.

Suppose that $\sigma \in \widetilde{\mathcal{S}}$ and fix $s_0 \in \mathbb{R}^* \setminus \{1\}$, $r_0 \in \mathbb{R}$ and $n \in \mathbb{N}$. Next choose $I \not\ni 1$ and $J \subset \mathbb{R}$ compact intervals such that $(s_0, r_0) \in I \times J$. By assumption, there exists a finite cover $\kappa \in \mathcal{V}(I \times J)$ such that for every $F \in \kappa$ there exists $\mathcal{O}_F \in \mathcal{I}$ with

$$\sigma(\mathcal{O}_F) > 1 - 1/n \quad \text{and} \quad \sigma_s * \delta_r(\mathcal{O}_F) < 1/n \quad \text{for all } (s,r) \in F.$$

Choosing $F \in \kappa$ for which $(s_0, r_0) \in F$ and applying (17) we have that σ and $\sigma_{s_0} * \delta_{r_0}$ are orthogonal, so $\sigma \in \mathcal{S}$.

It remains to show that $\mathcal{S}$ is dense. To this end we use the proof of Proposition 3.4 in [4]. Namely, in this proposition there is a construction of a weakly mixing flow $\mathcal{T}$ such that for a certain sequence of real numbers $u_k \to \infty$ we have: for each $l \in \mathbb{N}$

$$T_{-du_k} \to 10^{-l} \quad \text{for } d = 1 - 10^{-l} \text{ and} \tag{19}$$

$$T_{-cu_k} \to 0 \quad \text{uniformly in } c \in [1, 10^l] \tag{20}$$

(the convergence takes place in the weak operator topology). It follows that

$$\sigma_{\mathcal{T}_d} \perp \sigma_{\mathcal{T}_c} * \delta_t \tag{21}$$

for all $t \in \mathbb{R}$; indeed, (19) and (20) mean respectively

$$\xi_{u_k} \to 10^{-l} \quad \text{weakly in } L^2(\mathbb{R}, \sigma_{\mathcal{T}_d}),$$

and

$$\xi_{u_k} \to 0 \quad \text{weakly in } L^2(\mathbb{R}, \sigma_{\mathcal{T}_c}).$$

It is easy to see that the latter condition implies

$$\xi_{u_k} \to 0 \quad \text{weakly in } L^2(\mathbb{R}, \sigma_{\mathcal{T}_c} * \delta_t)$$

for each $t \in \mathbb{R}$, and the mutual singularity (21) follows.

Now, in view of (21), $\sigma_{\mathcal{T}} \perp \sigma_{\mathcal{T}_{c/d}} * \delta_{t/d}$, and since in (20) c can be replaced by $-c$, it follows that $\sigma_{\mathcal{T}} \in \mathcal{S}$. It is also clear that $\mathcal{S}$ is closed under taking absolutely continuous measures. Since $\operatorname{supp} \sigma_{\mathcal{T}} = \mathbb{R}$ [8], the result follows from (16). □

Recall also the following basic observation.

Lemma 17. *Let $\overline{s} = (s_j)_{j \geq 1}$ be a sequence of positive numbers and let $\overline{g} = (g_j)_{j \geq 1}$ be a sequence of uniformly bounded continuous functions. Then the set*

$$\mathcal{W}_{\overline{s},\overline{g}} = \left\{ \nu \in \mathcal{P}(\mathbb{R}) \colon \ (\exists\ t_n \to \infty)\, (\forall\ j \geq 1) \quad \xi_{s_j t_n} \to g_j \ \text{weakly in } L^2(\mathbb{R}, \nu) \right\}$$

is G_δ in $\mathcal{P}(\mathbb{R})$.

---

[8] This fact is well known for $\mathbb{Z}$-actions, e.g. [20], Chapter 3, and can be easily rewritten using special representation of flows. See also the proof of Theorem A in [22].

Proof. Let $(f_m)_{m\geq 1}$ be a sequence of continuous functions on $\mathbb{R}$ uniformly bounded by 1, which is linearly dense in $L^2(\mathbb{R},\nu)$ for every $\nu\in\mathcal{P}(\mathbb{R})$. Set

$$\mathcal{R}(n,\varepsilon)=\left\{\mu\in\mathcal{P}(\mathbb{R}):\sum_{m,j\geq 1}\frac{1}{2^{m+j}}\left|\int_{\mathbb{R}}\left(e^{2\pi i s_j n x}-g_j(x)\right)f_m(x)\,d\mu(x)\right|<\varepsilon\right\}.$$

The set $\mathcal{R}(n,\varepsilon)$ is open. To complete the proof it suffices to notice that

$$\mathcal{W}_{\bar{s},\bar{g}}=\bigcap_{\mathbb{Q}\ni\varepsilon>0}\bigcap_{m\geq 1}\bigcup_{n\geq m}\mathcal{R}(n,\varepsilon).$$

□

Lemma 18. *Let $H\subset\mathbb{R}_+^*$ be a countable multiplicative subgroup. Then for a typical $\nu\in\mathcal{P}(\mathbb{R})$ the measure $\eta:=\sum_{h\in H}a_h\nu_h$ (with $a_h>0$ and $\sum_{h\in H}a_h=1$) yields a Gaussian flow $\mathcal{T}^{\eta|_{\mathbb{R}_+}}$ with simple spectrum.*

Proof. Set $G=-H\cup H$ and let $H=\{s_i:\ i\geq 0\}$ ($s_0=1$). In [4], Danilenko and Ryzhikov constructed a rank-1 flow $\mathcal{T}$ preserving a σ-finite measure μ (the flow acts on $(X,\mathcal{B},\mu)$) such that if $\sigma=\sigma_{\mathcal{T}}$ denotes its maximal spectral type on $L^2(X,\mathcal{B},\mu)$ then the Gaussian flow

$$\mathcal{T}^{\left(\sum_{i\geq 1}\frac{1}{2^i}\sigma_{s_i}\right)|_{\mathbb{R}_+}}\text{ has simple spectrum.}\tag{22}$$

To prove this, they used the following properties of $\mathcal{T}$:

a) $T_{\sqrt{2}s}\in WCP(T_s)$ [9] for each $s\in H$,
b) $\frac{1}{q}I+\frac{q-1}{q}T_s\in WCP(T_s)$ for each $s\in H$ and $q\in\mathbb{N}$,
c) for each finite sequence $s_1<s_2<\dots<s_k$ of elements of H and each $1\leq l_0\leq k$ there exists $t_j\to\infty$ such that
  (i) $T_{t_js_j}\to\frac{1}{2k}I$ if $1\leq l\leq k, l\neq l_0$,
  (ii) $T_{t_js_{l_0}}\to\frac{1}{2k}T_{s_{l_0}}$.

Notice that the conditions a), b) and c) can be expressed as follows in terms of weak convergence of continuous and bounded functions in $L^2(\mathbb{R},\sigma)$:

a') for each $s\in H$ there exists a sequence $n_k\to\infty$ such that
$$\xi_{sn_k}\to\xi_{\sqrt{2}s},$$
b') for each $s\in H$ and $q\in\mathbb{N}$ there exists a sequence $n_k\to\infty$ such that
$$\xi_{sn_k}\to\frac{1}{q}+\frac{q-1}{q}\xi_s,$$
c') for each finite sequence $s_1<s_2<\dots<s_k$ of elements of H and each $1\leq l_0\leq k$ there exists $t_j\to\infty$ such that
  (i) $\xi_{t_js_j}\to\frac{1}{2k}$ if $1\leq l\leq k, l\neq l_0$,
  (ii) $\xi_{t_js_{l_0}}\to\frac{1}{2k}\xi_{s_{l_0}}$.

The arguments used in the proof of Theorem 4.4 in [4] show that for each continuous probability measure σ on $\mathbb{R}$ conditions a'), b') and c') imply the simplicity of spectrum of the flow $\mathcal{T}^{\left(\sum_{k\geq 1}\frac{1}{2^k}\sigma_{s_k}\right)|_{\mathbb{R}_+}}$. Moreover, by Lemma 17, the set of measures $\nu\in\mathcal{P}(\mathbb{R})$ satisfying these conditions is G_δ. We will show now that it is also dense in $\mathcal{P}(\mathbb{R})$. Notice that conditions a'), b') and c') hold also in $L^2(\mathbb{R},\nu)$ for any $\nu\ll\sigma$. Since $\sigma_{\mathcal{T}}$ is the maximal spectral type of a rank-1 infinite measure-preserving flow $\mathcal{T}$, the Gelfand spectrum of the corresponding Koopman representation is equal to $\mathbb{R}$. It follows that the topological support of $\sigma_{\mathcal{T}}$ is full and therefore the result follows from (16). □

---

[9] An operator Q belongs to the weak closure of powers WCP(R) if for an increasing sequence (m_j) of integers, $R^{m_j}\to P$ in the weak operator topology.

3. Proofs of theorems

Proof of Theorem 3. (based on Lemmas 12 and 16.) Using these two lemmas, for a "typical" (continuous, Kronecker) measure $\sigma \in \mathcal{P}([a,b])$ we have (with $a_h > 0$, and $\sum_{h\in H} a_h = 1$)

$$-H \cup H \subset I(\mathcal{T}^{\eta}),$$

where $\eta := \sum_{h\in H} a_h \sigma_h$ is a Kronecker measure and moreover

$$\sigma_s \perp \sigma * \delta_t \tag{23}$$

for each non-zero real $s \neq 1$ and arbitrary $t \in \mathbb{R}$. All we need to show is that when $s \notin -H \cup H$ then $\eta_s \not\equiv \eta$. However if $s \notin H$ then even more is true: $\eta \perp \eta_s * \delta_t$ for arbitrary $t \in \mathbb{R}$ and $s \notin \{0,1\}$. It follows that

$$\widetilde{\eta}_s \perp \widetilde{\eta} * \delta_t$$

for each $s \notin -H \cup H$ and $t \in \mathbb{R}$. In view of Theorem 9, it follows that $\mathcal{T}^{\eta}$ is disjoint from $\mathcal{T}^{\eta_s}$ (isomorphic to $\mathcal{T}^{\eta}_s$) for $s \notin -H \cup H$. In particular, $-H \cup H = I(\mathcal{T}^{\eta})$ and the result follows. ☐

Proof of Theorem 3. (based on Lemma 13.) Given $H \subset \mathbb{R}^*_+$ a multiplicative subgroup which is an additively $\mathbb{Q}$-independent set, in [6], there is a construction of a perfect compact set K such that $\widehat{K} := \bigcup_{h\in H} hK$ is independent and for $\widetilde{K} := -\widehat{K} \cup \widehat{K}$ the following holds: $(r\widetilde{K} + t) \cap \widetilde{K}$ is countable whenever $|r| \notin H$ and $t \in \mathbb{R}$ is arbitrary. Using Lemma 13 find a (continuous, Kronecker) measure $\sigma \in \mathcal{P}(K)$ such that $\eta := \sum_{h\in H} a_h \sigma_h$ is a Kronecker measure. Then η is concentrated on $\widehat{K}$. All we need to show is that if $|r| \notin H$, then the symmetrization of η_r is not equivalent to the symmetrization of η. This is however clear, since the symmetrization of η_r is a continuous measure concentrated on $r\widetilde{K}$. As in the previous proof we deduce that for $s \notin -H \cup H$ we obtain disjointness of the corresponding flows. ☐

Proof of Theorem 4. First notice that directly from Lemma 15, it follows that whenever σ is a Kronecker measure then for each $r_1, r_2 \in \mathbb{Q}^*$, $r_1 \neq r_2$, we have

$$\sigma_{r_1} \perp \sigma_{r_2} * \delta_t \quad \text{for each} \quad t \in \mathbb{R}.$$

It follows that $\widetilde{\sigma}_{r_1} \perp \widetilde{\sigma}_{r_2} * \delta_t$ for all $t \in \mathbb{R}$, so by Theorem 9, the Gaussian-Kronecker flows $\mathcal{T}^{\sigma_{r_1}}$ and $\mathcal{T}^{\sigma_{r_2}}$ are disjoint. In view of (6), it follows that $T_1^{\sigma_{r_1}} \perp T_1^{\sigma_{r_2}}$, thus $T^{\sigma}_{r_1} \perp T^{\sigma}_{r_2}$.

Now suppose that $T = T_\sigma : (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ is a Gaussian-Kronecker automorphism, i.e. $\sigma = \sigma_0 + \overline{\sigma}_0$ for a continuous Kronecker measure $\sigma_0 \in \mathcal{P}(\mathbb{T})$.[10] Denote by σ' the image of σ_0 via the map $\mathbb{T} \ni z \mapsto \operatorname{Arg}(z)/2\pi \in [0,1)$. Then σ' is a continuous Kronecker measure on $\mathbb{R}$ such that $(\xi_1)_*\widetilde{\sigma}' = \sigma$ and $\widetilde{\sigma}' * \delta_m \perp \widetilde{\sigma}'$ for all $m \in \mathbb{N}$. Denote by $\mathcal{H}$ the Gaussian space of the flow $\mathcal{T}^{\sigma'}$. Then the Koopman operator of $T_1^{\sigma'}$ has simple spectrum on $\mathcal{H}$ and its spectral type is $(\xi_1)_*\widetilde{\sigma}' = \sigma$, see Appendix in [16]. Since the spectral type of ζ_1 (with respect to $T_1^{\sigma'}$) is $(\xi_1)_*\widetilde{\sigma}' = \sigma$, it follows that $\zeta_1 \circ (T_1^{\sigma'})^n$, $n \in \mathbb{Z}$, span the space $\mathcal{H}$. Thus $T_1^{\sigma'}$ is isomorphic to T_σ. By the first assertion of the theorem, it follows that T^n_σ is disjoint from T^m_σ for any pair of distinct natural numbers.

In order to prove the second part of the theorem note that if s is irrational then the set $\{1, s\}$ is $\mathbb{Q}$-independent, so by Lemma 12 we can find a (continuous, Kronecker) measure $\sigma \in \mathcal{P}([a,b])$ such that $\eta := \frac{1}{2}(\sigma + \sigma_s)$ is a Kronecker measure. Since $\sigma_s \ll \eta$ and $\sigma_s \ll \eta_s$ the Gaussian-Kronecker flows $\mathcal{T}^{\eta}$ and $\mathcal{T}^{\eta_s}$ have a common non-trivial (Gaussian) factor. Its time one map is a common non-trivial factor of T_1^{η} and $T_1^{\eta_s}$ and it remains to notice that the Gaussian automorphism $T_1^{\eta_s}$ is isomorphic to T_s^{η}. ☐

---

[10] $\mathbb{T}$ stands for $\{z \in \mathbb{C} \colon |z| = 1\}$.

Proof of Theorem 5. Let $H = G \cap \mathbb{R}_+^*$ and let $(a_h)_{h\in H}$ be positive numbers such that $\sum_{h\in H} a_h = 1$. By Lemmas 16, 18 and Lemma 12 (applied to $H = \{1\}$) combined with Remark 2, there exists $\nu' \in \mathcal{P}_c(\mathbb{R})$ such that

(i) $\nu'_s \perp \nu' * \delta_t$ for all $s \in \mathbb{R}^* \setminus \{1\}$ and $t \in \mathbb{R}$;
(ii) the Gaussian flow $\mathcal{T}^{(\sum_{h\in H} a_h \nu'_s)|_{\mathbb{R}_+}}$ has simple spectrum
(iii) $\nu := \Delta(\nu') \in \mathcal{P}_c([a,b])$ is a Kronecker measure

(in fact, for a "typical" $\nu' \in \mathcal{P}_c(\mathbb{R})$ the properties (i)-(iii) hold). Since the conditions (i) and (ii) hold also for any measure absolutely continuous with respect to ν, the Kronecker measure ν satisfies (i) and (ii) as well. Therefore, setting $\sigma := \sum_{h\in H} a_h \nu_s$, by (ii), the Gaussian flow $\mathcal{T}^{\sigma}$ has simple spectrum. The same argument as in the proof of Theorem 3 shows that (i) together with (ii) imply $I(\mathcal{T}^{\sigma}) = -H \cup H$ and $\mathcal{T}^{\nu_s} \perp \mathcal{T}^{\nu_r}$ whenever $|r| \neq |s|$. Each Kronecker measure ν_h, $h \in H$ is an FS measure so, by Proposition 10, it follows that $\sigma = \sum_{h\in H} a_h \nu_s$ is an FS measure [11], which completes the proof. ☐

Proof of Theorem 6. The first part follows from Lemma 16 along the same lines as the first proof of Theorem 3 (for $H = \{1\}$).

In view of Corollary 2 in [16], a typical flow $\mathcal{T}$ has the SC property,[12] which is equivalent to the fact that $\mathcal{T}^{\sigma_{\mathcal{T}}}$ has simple spectrum. In particular, it implies that $\mathcal{T}^{\sigma_{\mathcal{T}}}$ is GAG.

In order to prove that $\sigma_{\mathcal{T}} \perp (\sigma_{\mathcal{T}})_s * \delta_r$, $s \in \mathbb{R}^* \setminus \{1\}$, $r \in \mathbb{R}$ for a typical flow $\mathcal{T}$ we follow the proof of Theorem 3.2 from [4] (using Lemma 16 and the existence of a flow satisfying (21)). Since $\mathcal{T}^{\sigma_{\mathcal{T}}}$ is GAG for a typical flow $\mathcal{T}$, by Proposition 9, it follows that $\mathcal{T}^{(\sigma_{\mathcal{T}})_s}$ and $\mathcal{T}^{(\sigma_{\mathcal{T}})_r}$ are disjoint wherever $|r| \neq |s|$. ☐

Question. Is there a Kronecker measure $\sigma \in \mathcal{P}(\mathbb{R}_+)$ such that $I(\mathcal{T}^{\sigma})$ is uncountable?
This question is to be compared with Ryzhikov's question whether there is a weakly mixing, non-mixing flows with uncountable group of self-similarities, see [3], Problem (1).

References

[1] J. Bourgain, P. Sarnak, T. Ziegler, *Disjointness of Möbius from horocycles flows*, arXiv 1110.0992v1.
[2] I.P. Cornfeld, S.V. Fomin, Y.G. Sinai, *Ergodic Theory*, Springer-Verlag, New York, 1982.
[3] A. Danilenko, *Flows with uncountable but meager group of self-similarities*, to appear in Contemporary Math.
[4] A. Danilenko, V.V. Ryzhikov, *On self-similarities of ergodic flows*, to appear in Proc. London Math. Soc.
[5] C. Foiaş, S. Stratila, *Ensembles de Kronecker dans la théorie ergodique*, C.R. Acad. Sci. Paris, série A **267**, 166-168.
[6] K. Frączek, M. Lemańczyk, *On the self-similarity problem for ergodic flows*, Proc. London Math. Soc. **99** (2009), 658-696.
[7] H. Furstenberg, *Disjointness in ergodic theory, minimal sets and diophantine approximation*, Math. Syst. Th. **1** (1967), 1-49.
[8] H. Furstenberg, *Recurrence in Ergodic Theory and Combinatorial Number Theory*, Princeton University Press, Princeton, New Jersey, 1981.

---

[11] We use here the elementary fact that the L^2-limit of a sequence of Gausian variables remains Gaussian.

[12] The SC property means that if we set $\sigma = \sigma_{\mathcal{T}}$ then for each $n \geq 2$ the conditional measures of the disintegration of $\sigma^{\otimes n}$ over σ^{*n} via the map $\mathbb{R}^n \ni (x_1, \dots, x_n) \mapsto x_1 + \dots + x_n \in \mathbb{R}$ are purely atomic with $n!$ atoms.

[9] E. Glasner, *Ergodic Theory via Joinings*, Mathematical Surveys and Monographs **101**, AMS, Providence, RI, 2003.
[10] A. del Junco, D. Rudolph, *On ergodic actions whose self-joinings are graphs*, Ergodic Theory Dynam. Systems **7** (1987), 531-557.
[11] A. Katok, J.-P. Thouvenot, *Spectral Properties and Combinatorial Constructions in Ergodic Theory*, Handbook of dynamical systems. Vol. 1B, 649–743, Elsevier B. V., Amsterdam, 2006.
[12] T.-W. Körner, *Some results on Kronecker, Dirichlet and Helson sets*, Annales Inst. Fourier, **20** (1970), 219-324.
[13] J. Kułaga, *On the self-similarity problem for smooth flows on orientable surfaces*, Ergodic Theory Dynam. Systems (2011) (published online).
[14] M. Lemańczyk, *Spectral Theory of Dynamical Systems*, Encyclopedia of Complexity and System Science, Springer-Verlag (2009), 8554-8575.
[15] M. Lemańczyk, F. Parreau, *On the disjointness problem for Gaussian automorphisms*, Proc. Amer. Math. Soc. **127** (1999), 2073-2081.
[16] M. Lemańczyk, F. Parreau, *Special flows over irrational rotations with the simple convolutions property*, preprint available `http://www-users.mat.umk.pl/~mlem/publications.php`
[17] M. Lemańczyk, F. Parreau, J.-P. Thouvenot, *Gaussian automorphisms whose ergodic self-joinings are Gaussian*, Fund. Mah. **164** (2000), 253-293.
[18] L.-A. Lindahl, F. Poulsen, *Thin Sets in Harmnonic Analysis*, Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, Inc. New York, 1971.
[19] B. Marcus, *The horocycle flow is mixing of all orders*, Invent. Math. **46** (1978), 201-209.
[20] M.G. Nadkarni, *Spectral Theory of Dynamical Systems*, Birkhäuser Advanced Texts 1998.
[21] F. Parreau, *On the Foiaş and Stratila theorem*, Proc. Conference on Erg odic Theory and Dynamical Systems, Toruń 2000, 106-108, available `http://www-users.mat.umk.pl/~mlem`
[22] W.C. Ridge, *Spectrum of a composition operator*, Proc. Amer. Math. Soc. **37**, (1973), 121-127.
[23] T. de la Rue, *Joinings in ergodic theory*, Encyclopedia of Complexity and System Science, Springer-Verlag (2009), 5037-5051.
[24] V.V. Ryzhikov, *Intertwinings of tensor products, and the centralizer of dynamical systems*, Sb. Math. **188** (1997), 67-94.
[25] V.V. Ryzhikov, *On disjointness of mixing rank one actions*, arXiv:1109.0671v1

Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, ul. Chopina 12/18, 87-100 Toruń, Poland
E-mail address: `fraczek@mat.umk.pl, joanna.kulaga@gmail.com, mlem@mat.umk.pl`